

Results of Examining Small Impact Crater Populations on the Moon

Michelle Kirchoff, Clark Chapman, Simone
Marchi, William Bottke, Brian Enke, and
Kristen Sherman

Southwest Research Institute
Center for Lunar Origin and Evolution




NASA
LUNAR SCIENCE
FORUM

July 18, 2012

NASA
LUNAR SCIENCE
INSTITUTE



GOAL: INVESTIGATE LUNAR IMPACTORS' SIZE-FREQUENCY DISTRIBUTIONS (SFDs) THROUGH TIME AND IMPROVE ESTIMATES OF IMPACT RATES SINCE THE LHB




Motivation

- » Previous work needs updating
e.g., Wilhelms et al. (1978), Hartmann et al. (1981), Neukum & Ivanov (1994)
- » Little work for this age & diameter range
- » Outstanding LROC imaging - WAC global mosaic
<http://lroc.sese.asu.edu/>
- » Recent developments in numerical approaches
e.g., Bottke et al. (2000, 2002, 2005), Marchi et al. (2009)

**OUR APPROACH: COMPILING AND
ANALYZING SMALL, SUPERPOSED
CRATER SFDs ON THE FLOORS OF
SEVERAL LARGE LUNAR CRATERS**



Selected Mid-Sized Craters



» Wide distribution in latitude and longitude, and age

Compilation & Analysis

- » Measure small superposed craters and assign degradation class (1-4) & obvious secondary (OS; clusters/chains) in JMARS
- » Compute model ages with Marchi et al. (2009) Model Production Function (MPF)
 - » Compare small superposed crater SFDs to MPF
 - » Assign Stöffler et al. (2006) epoch & compare to past USGS identifications




Stöffler et al. (2006)


Proposed time periods
for lunar time units

Time unit	Time (Gyr) options	Time (Gyr) adopted
Copernican Period	a) 2.1 - 0 b) 1.0 - 0 c) 0.8 - 0	0.8 - 0
Eratosthenian Period	a) 3.2 - 2.1 b) 3.2 - 1.0 c) 3.2 - 0.8	3.2 - 0.8
Late Imbrian Period	a) 3.75 - 3.2 b) 3.72 - 3.2	3.75 - 3.2
Early Imbrian Period	a) 3.85 - 3.72 b) 3.85 - 3.75 c) 3.77 - 3.72 d) 3.77 - 3.75	a) 3.85 - 3.75 b) 3.77 - 3.75
Nectarian Period	a) 3.92 - 3.85 b) 3.92 - 3.77 c) 3.85 - 3.77	a) 3.92 - 3.85 b) 3.92 - 3.77
Pre- Nectarian Period	a) 4.52 - 3.92 b) 4.52 - 3.85	4.52 - 3.92

36/40 SSC SFDs are consistent with MPF



Crater Floor Model Ages



» Indicates LHB possibly ended with extended tail & little cratering < 2 Ga

Crater Floor (CF) Model Ages

Crater Floor	Model Age	Associated Epoch (Stöffler et al., 2006)	Previous USGS Epoch
Vavilov	1.7 ± 0.1	Eratosthenian	Copernican
Hayn	1.8 ± 0.8	Eratosthenian	Copernican
Theophilus	3.0 ± 1.2	Eratosthenian	Copernican
Geminus	3.2 ± 0.7	Eratos./Imbr.	Eratosthenian
Hausen	3.5 ± 0.2	Late Imbrian	Eratosthenian
Robertson	3.7 ± 0.1	Late Imbrian	Copernican
Birkeland	3.8 ± 0.1	Early Imbrian	Eratosthenian
Lobachevskiy	3.8 ± 0.1	Early Imbrian	Late Imbrian
Hahn	3.8 ± 0.1	Early Imbrian	Late Imbrian
Arnold	3.8 ± 0.1	Early Imbrian	Late Imbrian
Baillaud	3.9 ± 0.1	Nectarian	Late Imbrian
Piccolomini	3.9 ± 0.1	Nectarian	Imbrian
Bridgman	3.9 ± 0.1	Nectarian	Imbrian
Vlacq	3.9 ± 0.1	Nectarian	Imbrian
Ansgarius	3.9 ± 0.1	Nectarian	Imbrian
Joule	4.0 ± 0.1	Pre-Nectarian	Imbr./Nect.
Laue	4.0 ± 0.1	Pre-Nectarian	Imbrian
Freundlich	4.0 ± 0.1	Pre-Nectarian	Imbrian
Neumayer	4.0 ± 0.1	Pre-Nectarian	Nectarian






»For ~50% of CFs, our ages are older than previous

»Includes all formerly Copernican & Eratosthenian crater floors

»Issues with our data/calculations?


»More interesting: issues w/ USGS interpretations and CFs are actually older?

4/40 SSC SFDs NOT consistent w/ MPF




MPF Inputs Adjusted or SFD Geologically Modified?

Impactor





Population 1
Strom et al. (2001)

Scaling Law




Rosenberger
w/ Layered Scaling Law
Marchi et al. (2011)

Resurfacing



Obvious Secondaries



- » Wide variation in obvious secondary SFD slopes
- » Related to formation of individual secondary crater fields?

Conclusions


- » Agreement between most of SSC SFDs and MPF further validates MPF & indicates SSC SFDs are likely largely characterized by primary craters
- » Model ages of original CFs indicate LHB had an extended tail
- » Ages older than initially ascertained => may be less cratering during Copernican/Early Eratosthenian (< 2 Ga) than originally thought
- » Few SSC SFDs not consistent with MPF => Likely modified by geologic activity, but MPF inputs can be adjusted to fit; is this appropriate?
- » Obvious secondary SFDs have widely varying slopes that likely are representative of different types of secondary cratering

Acknowledgments: NLSI for funding support, JMARS support team, and my co-authors

SSC SFD Unchanging?

$D < 3 \text{ km}$
Yes

=> Impactor SFD
slope remain stable
over 1-4 Ga?




$D > 3 \text{ km}$
??


Poor statistics:
approaching size of
counting areas

Trend: transition to
shallower slopes??

Counting Areas



Analysis



Stöffler et al. (2006)
Proposed time periods
for lunar time units

Time unit	Time (Gyr) options	Time (Gyr) adopted
Copernican Period	a) 2.1 - 0 b) 1.0 - 0 c) 0.8 - 0	0.8 - 0
Eratosthenian Period	a) 3.2 - 2.1 b) 3.2 - 1.0 c) 3.2 - 0.8	3.2 - 0.8
Late Imbrian Period	a) 3.75 - 3.2 b) 3.72 - 3.2	3.75 - 3.2
Early Imbrian Period	a) 3.85 - 3.72 b) 3.85 - 3.75 c) 3.77 - 3.72 d) 3.77 - 3.75	a) 3.85 - 3.75 b) 3.77 - 3.75
Nectarian Period	a) 3.92 - 3.85 b) 3.92 - 3.77 c) 3.85 - 3.77	a) 3.92 - 3.85 b) 3.92 - 3.77
Pre-Nectarian Period	a) 4.52 - 3.92 b) 4.52 - 3.85	4.52 - 3.92




SFD Interpretations - Age

Crater	Relative Age	Absolute Age	Stöffler Epoch	USGS Epoch
Hayn	1800 ± 1000	1.1 ± 0.2	Cop./Erat.	Copernican
Vavilov	3600 ± 2100	1.3 ± 0.2	Cop./Erat.	Copernican
Aristoteles	6200 ± 1800	3.5 ± 0.2	Late Imbrian	Eratosthenian
Theophilus	7100 ± 2000	3.4 ± 0.2	Late Imbrian	Copernican
Hale	7400 ± 2800	3.6 ± 0.1	Late Imbrian	Late Imbrian
Hausen	7600 ± 900	3.6 ± 0.1	Late Imbrian	Eratosthenian
Lyman	8300 ± 2400	3.6 ± 0.1	Late Imbrian	Late Imbrian
La Pérouse	10000 ± 2800	3.7 ± 0.1	Late Imbrian	Late Imbrian
Roberts	13600 ± 2500	3.8 ± 0.2	Early Imb./Nect.	Early Imb./Nect.
Langmuir	15500 ± 3900	3.8 ± 0.1	Early Imb./Nect.	Early Imbrian
Arnold	16000 ± 1900	3.8 ± 0.1	Early Imb./Nect.	Early Imbrian
Bose	16500 ± 2500	3.8 ± 0.2	Early Imb./Nect.	Early Imbrian
Lobachevskiy	17200 ± 3100	3.9 ± 0.1	Nectarian	Late Imbrian
<u>Birkeland</u>	<u>18800 ± 4200</u>	<u>3.9 ± 0.2</u>	<u>Nectarian</u>	<u>Eratosthenian</u>
<u>Hahn</u>	<u>19400 ± 5400</u>	<u>3.9 ± 0.1</u>	<u>Nectarian</u>	<u>Late Imbrian</u>
Al-Biruni	19800 ± 3300	3.8 ± 0.1	Early Imb./Nect.	Early Imbrian
Laue	25000 ± 3900	3.9 ± 0.2	Nectarian	Early Imbrian
Baillaud	25600 ± 2300	3.9 ± 0.1	Nectarian	Early Imbrian
<u>Millikan</u>	<u>25800 ± 3100</u>	<u>4.0 ± 0.1</u>	<u>Pre-Nectarian</u>	<u>Early Imb./Nect.</u>
Freundlich	26300 ± 3800	3.9 ± 0.1	Nectarian	Early Imbrian

» SFDs that match MPF (black) have wide age distribution

» SFDs that are shallower than MPF (red) all relatively old

SFD Interpretations = Distribution



- » SFDs that match MPF (yellow) appear widely distributed in latitude & longitude
- » Majority of SFDs that are shallower than MPF (red) on farside – pattern??

